

## เอกสารอ้างอิง

- บัญญัติ สุขศรีงาม. 2532. จุลชีววิทยาทั่วไป. สำนักพิมพ์โอเดียนสโตร์ กรุงเทพฯ.
- สรณรัชฎ์ กาญจนวณิชย์และ สตีเฟน ทิลลิง. (ไม่ระบุปีที่ตีพิมพ์). คู่มือจำแนกพันธุ์สัตว์ไม่มีกระดูกสันหลังในบึงและลำธารไทย. มุลนิธิโลกสีเขียว & Field Studies Concl.
- Benavides M.S. 1994. Algal Periphyton in Two Rivers in Costa Rica with Special Reference to Diatoms Organic Pollution and Altitudinal Differentiation. Ph.D. Thesis. Institute of Botany, Innsbruck University, Austria.
- Branco C.C.Z., Necchi O.Jr., Branco L.H.Z. 2001. Taxonomy of Chaetophoraceae (Chaetophorales, Chlorophyta) in Lotic Ecosystem from Sao Paulo State, South-Eastern Brazil. Journal of Phycologia, 40(4):420.
- Flint L.H. 1960. Freshwater Red Algae of North America. Vantage Press, New York.
- Gardarsky A. 1986. *Cladophora moravica* (Drorak) Comb Nora and *C. basiramosa* Schmidle Two Interesting Rivewine Species of the Green Filamentous Algae from Czechoslovakia. Arch. Hydrobiol, 73:49-77.
- Kunpradid, T. and Y. Peerapornpisal. 2002. Using of Macroalgae Distribution as Biomonitor in Ping River. Journal of Science, Chulalongkorn University.
- Kunpradid, T., Wannathong P., Leeahakhangkri P. and Y. Peerapornpisal . 2004. Benthic Algae as Biomonitor to Assess the Water Quality in Ping Watershed, Northern Thailand. Societies Internationals Limnology Congress, University of Helsinki, Lahti, Finland, 8-14 August 2004
- Mustow, S.E. (2002) Biological Monitoring of Rivers in Thailand : Use and Adaptation of the BMWP Score. Netherlands Journal Hydrobiologia 479 : 191-229.
- Necchi O., Pascoaloto D. 1995. Morphometry of *Compsopogon coeruleus* (Compsopogonaceae, Rhodophyta) Populations in a Tropical River Basin of Southeastern Brazil. Journal of Phycologia, 32:150-168.
- Palmer R.G. 1970. Algae and Water Pollution. London Press, London.
- Peerapornpisal Y., Pekthong T., Waiyaka P., Promkutkaew S. 2000. Diversity of Phytoplankton and Benthic Algae in Mae Sa Stream, Doi Suthep-Puk National Park, Thailand. Journal of Natural History Bulletin of the Siam Society, 48:193-211.
- Peerapornpisal Y., Kunpradid T. and Suphan S. 2003. Benthic Diatoms and Macroalgae as the Biomonitors to Assess the Water Quality in Mekong River. Report on the Development of Ecological Health Monitoring Program of Mekong River Commission.

- Sheath R.G., Cole C.M. 1992. Use of *Cladophora glomerata* to Monitor Heavy Metals in Rivers. *Journal of Applied Phycology*, Newyork.
- Walley W.J., Hawkes H.A. 1997. A Computer-Based Development of the Biological Monitoring Working Party Score System incorporating Abundance Rating, Site Type and Indicator Value. *Wat. Res.*, 31(2):201-210.
- Walley W.J., Fontama V.N. 1998. Neural Network Predictors of Average Score Per Taxon and Number of Families at Unpolluted River Sites in Great Britain. *Wat. Res.*, 32(3):613-622.
- Whitton , B.A. and G. K. Martyn. 1995. Use of Algae and Other Plants for Monitoring Rivers. *Australian Journal of Ecology* , 20 : 45 - 56.
- Wong S.L., Nakamoto L., Wainwright J.F. 1997. Detection of Toxic Organometallic Complexes in Wastewater Using Algal Assays. *Arch. of Enivonmental Contamination & Toxicology*, 32(4):358-366.
- Tada M. 2002. *The Fresh-Water Algae*. W.M.C. Brown Company Publishers, Iowa.

